

IEGSEUI is a registered trademant of Akcea Therspeutics, Inc., discovered and developed by lonis ₱harmaceuticals and distributed by 50bi, Inc. © 2024. Akcea Therspeutics, Inc. All rights reserved. REF-11441 XX/24

TEGSEDI® PATIENT WALLET CARD

Carry this card with you at all times. SHOW THIS CARD if you go to the emergency room or see any healthcare provider.

Tell any healthcare provider that sees you that you are being treated with TEGSEDI.

Prescriber's name (first & last):	
	- (4)
Prescriber phone #:	Tegsedi [®]
	— (inotersen) injection

Patient information

TEGSEDI may cause side effects that are severe or life threatening at any time during treatment.

Call your doctor or go to the emergency room RIGHT AWAY if you have any of these symptoms:

- Unusual bruising or a rash oftiny reddish-purple spots, often on the lower legs
- Bleeding from skin cuts that does not stop or oozes
- Bleeding from your gums or nose
- Blood in urine or stools
- Bleeding into the whites of your eyes
- Sudden severe headaches or neck stiffness
- Vomiting blood or coughing

- up blood
- Abnormal or heavy menstrual periods
- Puffiness or swelling in your face, feet, or hands
- New onset or worsening shortness of breath and coughing
- Bloodinyoururine or brownurine
- · Foamy urine (proteinuria)
- Passing less urine than usual

Important information for healthcare providers

This patient is receiving TEGSEDI for the treatment of the polyneuropathy of hereditary transthyretin-mediated amyloidosis in adults.

TEGSEDI has a risk of serious bleeding with severe thrombocytopenia and a risk of glomerulonephritis. Patients treated with TEGSEDI are being regularly monitored for these risks via the following lab results: platelets, estimated glomerular filtration rate (eGFR), urinalysis, and urine protein to creatinine ratio (UPCR).